



SYRACUSE ENDOSCOPY ASSOCIATES, LLC

5000 CAMPUSWOOD DRIVE, EAST SYRACUSE, NY 13057

WWW.SYRACUSEGASTRO.COM

(315) 234-6677

Medications that need to be held prior to procedures


DO NOT TAKE 7 DAYS PRIOR:

Phentermine (Adipex-P, Suprenza)
 Phentermine/Topiramate (Qsymia)
 Orlistat (Xenical, Alli)
 Lorcaserin (Belviq)
 Diethylpropion
 Methamphetamine (Desoxyn)
 Liraglutide (Saxenda)
 Benzphetamine (Didrex, Regimex)
 Phendimetrazine (Bontril PDM)
 Carafate
 Iron
 Multivitamins
 Herbal meds
 Fish oil
 Vitamins A&E

DO NOT TAKE 2 DAYS PRIOR:

Lisdexamfetamine (Vyvanse)
 Dextroamphetamine & Amphetamine (Adderall)
 Any ADHD medications
 Sildenafil (Viagra)
 Tadalafil (Cialis)

 No Marijuana or recreational drugs



If you are a diabetic patient, please call the doctor that manages your diabetes and let them know you will be prepping for a colonoscopy.

Advise them of the medication changes you need to make and work with your doctor on how to manage your blood sugar during your prep time.

DIABETIC AND WEIGHT LOSS MEDICATIONS

The following medications **MUST BE STOPPED** as instructed below

GLP-1 MEDICATIONS

- If you are taking a GLP-1 medication on a **DAILY BASIS** you will need to skip your dose the day of the procedure.
- If you are taking a GLP-1 medication on a **WEEKLY BASIS** you will need to skip your dose the week before your procedure.

Trulicity (Dulaglutide)
 Bydureon BCise or Byetta (Exenatide)
 Ozempic, Wegovy or Rybelsus (Semaglutide)
 Saxenda or Victoza (Liraglutide)
 Adlyxin or Lyxumia (Lixisenatide)
 Mounjaro or Zepbound (Tirzepatide)
 Soliqua (Insulin glargine/Lixisenatide)
 Beinaglutide
 PEG-loxenatide

SGLT2 MEDICATIONS

- Hold these medications for 3 days

Jardiance (Empagliflozin)
 Brenzavvy (Bexagliflozin)
 Invokana (Canagliflozin)
 Farxiga or Xigduo (Dapagliflozin)
 Inpefa (Sotagliflozin)
 Suglat (Ipragliflozin)
 Invokamet (Canagliflozin plus Metformin HCl)
 Synjardy (Empagliflozin plus Metformin HCl)
 Qtern (Dapagliflozin and Saxagliptin)

 Steglatro (Ertugliflozin) • Hold this medication for **4 days**